PROTOCOL DREDGE

Cruise No. SO 37	Stat. No. Equipm. 5 DK	Location SO-37/1	Area H Tiefsee
Date 18.05 1985	Protocol written by : Halbach / Herzig		

Pinger on wire 300 m from dredge

Tº Siehe tension recorder

	Time GMT	Latit. N S	Longit. W	Water depth	Wire length	Tension (t)	~~Remarks~~ Kurs
begin operation	22.45	23°04	176°04'08	2100	0	0	
begin lowering	22.48	23°04	176°04'09	2100	0	1	
+1. bottom contact	23.23	23°02	176°04'12	2107	2166	2	
	23.31	22°84	176°04'13	1877	2250	2	
Bemerkungen: -	23.35	22°74	176°04'14	1895	2214	2	
	23.49	22°50	176°04'00	2150	2214	5.5	320°
Dredge Boden.	23.56	22°37	176°04'09	2150	2214	6.0	280°
	00.00	22°29	176°04'22	2120	2361	2.0	308°
Kontakt von	00.07	22°17	176°04'38	2100	2361	6.0	280°
2100 - 1400 m*	00.19	22°07	176°04'70	1900	2461	3.0	286°.
	00.24	22°04	176°04'76	1800	2511	3.0	286°
*edle Tiefe	00.47	21°94	176°05'16	1668	2501	4.6	287°
	00.53	21°93	176°05'20	1600	2506	7.1	310°
	00.57	21.93	176°05'23	1500	2466	8.0	281°
	01.00	21.92	176°05'25	1500	2454	8.5	271°
	01.10	21.92	176°05'26	1400	2434	8.7	293°
	01.31	21.89	176°05'34	1350	2205	5.2	293°
	01.41	21.88	176°05'35	1300	2033	7.1	338°
	01.50	21.93	176°05'28	1300	1600	8.2	
begin heaving	01.45	21.92	176°05'29	1300	1780	3.5	
end heaving	02.22	21.83	176°05'50			1.8	
end operation	02.34	21.79	176°05'73				

+Corrected data 0° 23.26' N 176 4.39' W

ship course ..... calcul. mean water depth 1750 m

total time of bottom contact 147 min ship velocity 0.5- 1.5 kts

bottom topography steiler Hang length of dredge section 1:2 m

Schlepp

Result

250 kg Total (60 auf 400 kg - Gesteine!!) 150 kg Substrate

.... kg Nodules 60 kg Sediment (Konsolidiert ?)

5-13

SO32 5DK

	GMT	Lat S	Long W	Depth	Wire Out
Begin operation:	22.45	0°23'04	176°04'00	2100m	0
Lowering:	22.48	0°23'05"	176°04'09"	2100m	0
On bottom contact	23.23	0°23'02"	176°05'12"	2107m	2165
Begin heaving:	01.45	0°21'32"	176°05'20"	1300-1780	3.5
End heaving:	02.22	0°21'83"	176°05'-3"		1.3
End operation:	02.34	0°21'79"	176°05'77"		
[illegible]		0°23'26" S,	176°4'39" W		
[illegible]		1750m			

Corrected bearings: 0°23'26"S, 176°4'39"W
Corrected Bathy: 1750m

250 kg total wt.

% talus / % outcrop in 75-30% bottom fraction

% crusts in total dredge: ~ 90% 0.5 mm or more

0% nodules

{ thickest crust = 3 cm
thinnest crust = 5 mm } 0.30 range

sulphide topography: plateau [illegible]

Oxic generation crust [illegible]

grey / brown: Yellowish brown [illegible]

	1	2	3
Rock Type:	Alkali Olivine Basalt	→ do - but w/ Pyroxene	(?) do-, Aphyric, Scoriaceous
Color:	medium dark gray (N4) to Grayish-green (5YR3/1) to medium olive gray (5Y5/1)	Dark gray (N3) to Pale yellowish brown (10YR 6/2)	→ do-
Texture:	Intergranular	→ -do-	→ -do-
Phenocrysts: (Single pcts)	Olivine, ±5 mm dia, ±3% Range: 1-15 mm dia.	→ - do- ± 3 mm dia, ± 5%	Aphyric (rare augite)
		1-5 mm dia.	
		Pyroxene (augite?) ±10%, ± 3 mm dia.	
		→ do - Range: 1-10 mm dia	
Groundmass:	Olivine, Plagioclase, opaques, Pyroxene } not determined	Olivine (much less than 5% pl.), Plagioclase, Pyroxene, opaques } not determined	} - do - ? (beyond resolution)
Vesicles:	% absent (dense) - rare irregular ±5 mm dia vugs.	→ do absent (dense)	30-50%
	filling: empty	→ 2 cm dia	round-oval, ± 1 mm dia
		phillipsite, calcite	empty, rarely calcite
Alteration:	extensive (smectitization, oxidation, hydration, carbonation)	superficial-moderate (smectitization of olivines)	extensive (smectitization, oxidation, zeolitization(?) cross-cut by numerous calcite veinlets)
Remarks:	pebble to boulder-size angular-to-subangular clasts set in several different volcanogenic breccia from this site; primary outcrop probably not exposed; Olivine pxts are generally individual and are skeletonized by smectitization and carbonation (smectite is pale yellow-green, probably saponite); rock is highly fractured, bearing stains of $Fe(OH)_3$ (?) & MnO_2 along the fracture faces; may have all been erupted subaqueously as pillow basalt (numerous pieces bear several mm-thick palagonite zones which grade directly to adhering hyaloclastite; serves as substrate for ±5 mm thick deposits of knobby, wormy single generation Mn-oxide (dendritic type)	→ -do-	→ -do-
		subangular - subrounded	angular
		→ -do-	→ -do-
		→ -do-	→ -do-
		→ -do-	
		iddingsite	
			→ -do-
		contact sharp w/ V.B.'s or presents a weathered laterized upper(?) surface	
		- freshest basalt, this dredge -	→ -do-
			most pieces appear to have been A a - type flows since the lavas are scoriaceous & include fragments of other vesicular - to scoriaceous bas

← may all be related to one volcanic edifice

4 5 6

Rock Type: Andesite (Hawaiite) → -do- → -do-

Color: (Olive gray 5Y 5/1) Medium gray (N5) Light olive gray (5Y 6/1)

Texture: Intergranular → -do- → -do-

Phenocrysts (single pxls) enclosed: Hornblende (?): ≤ 3 mm dia. avg. 10%; Augite (?): ≤ 1 mm dia. avg. 5%; Plagioclase ≤ 1 mm dia. avg. 3%; Olivine (?): ≤ 1 mm dia. avg. 2%

Groundmass: 5%, ≤ 1 mm dia. avge; 5%, ≤ 1 mm dia. avge; +%, ≤ 1 mm dia. avge; 1%, ≤ 1 mm dia. avge

Groundmass: Hornblende, Augite, Plagioclase, Olivine, Opaques — not determined → -do- → -do-

Secondary: % altered (dense) → -do- → -do-

Vesicles: moderate (Oxidation, Silicification, Zeolitization) → Slight (oxidation, silicification) → extensive (Silicification, Oxidation)

Lithic fragments: fine-to-boulder size → -do- → -do-

Chilled vitric clasts set in hornblende/augite hyaloclastite matrix → -do- → -do-

Glass appears as resistant needles and lathes with adamantine luster, angular cleavages, in quite random orientation in the groundmass; plagioclase occurs as small laths or equidimensional prisms; augite occurs as clusters of stubby eight-sided crystals with rt.-angle cleavages; olivine is rare and rarely altered to whitish $CaCO_3$ and skeletonized; exposed (to seawater) surfaces coated with $Fe(OH)_3$ (?) which does not serve as substratum for deposit of MnO_2.

Subrounded → angular

-do- → -do-

somewhat flow oriented → highly flow-oriented

-do- → -do-

-do- → -do-

-do- → -do-

intrusive? (trap?)

All have adhering hyaloclastite

7 8 9

ock Type: Andesitic? Hyaloclastite → -do- → -do-

Moderate yellowish-brown (10YR 5/4) (mean color); dividual units quite variable: olorful brown (5YR 4/4) to light rown (5YR 5/6) to Moderate yellowish-rown (10YR 5/4) to Dark yellowish-range (10YR 6/6) → -do- ranges from Olive gray (5Y 3/1) to light olive-gray (5Y 6/1) to moderate brown (5YR 4/4) to dark yellowish-brown (10YR 4/2) to brown (5YR 5/6) to Moderate yellowish-brown (10YR 5/4) to dark yellowish-orange (10YR 6/6) ≠ yellowish-olive-gray to (mean color 5Y 6/2) grayish-orange (10YR 7/4) impression of ... weathering rinds is the most ... fragments ... zeolites ... Fe(OH)

ture: Pyroclastic → -do- → -do-

mework rains: Cinder-size andesitic frit (±glass) clasts; -do- also includes fragments (pebbles gravel) of dense andesite ≠ generally

ent: Calcite, zeolites ?; ±20% → -do- → -do-, but much less abundant

ange (mean Size ±) mm → -do- → -do-

ape Grain Shape: subangular → -do- ≠ Subrounded

ting: moderately poorly-sorted ≠ Very poorly-sorted ≠ Moderately poorly sorted

sity: Extremely high → -do- → -do-

sibility: Moderately low → -do- ≠ Very high

n: Total (Silicification, Zeolitization, Oxidation, Carbonation, hydration) ≠ nearly complete except for a few fresher included clasts ≠ incipient to moderate → -do- → -do-

rks: Appears to be the most ... of several varieties recovered by dredging at the ... station: composed of andesitic ... altered, glassy glass fragments. Much of the rock is open space (interstices) ... superficially with calcite and zeolites, enough to bind the matrix together. oxidation is the pronounced alteration, although the very abundant hornblende euhedra (±5%) seem unaltered: serves as substrate for patinate MnO_2 deposit & $Fe(OH)_3$?

→ similar to 5DK7 but with fresher cinders and rock fragments → -do-, but with more fresher material

-do- probably represents the source of dense andesites (hawaiites? mugearites?) found in dredge

→ ? (not observed on my samples) ≠ Serves as substrate for smoothed, warty (botryoidal) veneer-thin (0.5 mm) MnO_2 deposits

Sample 108:	5DK 10	5DK 11	5DK 12/13
Rock Type:	AOB(?) Hyaloclastite →	-do- (prob. related to 5DK2) →	-do- ("Matrix Cement" for V.B. 5DK13)
Color:	Light yellow (5Y 8/6) (mean color) to Dark reddish-brown (10R3/6) (lapanite casts) to Moderate reddish-brown (10R4/6) to Light brown (5YR 6/4) →	-do- (larger clasts) →	grayish-yellow (5Y 8/4) (cement) (mean color) To light brown (5YR 4/4) (matrix clasts)
Texture:	Pyroclastic →	-do- →	-do-
Framework Grains:	ash-Cinder-Size AOB frit ~40% →	Cinder frit, 40-60%. avge size, ~½mm	± 25%, ~ 5mm avg dia.
Cement:	Nontronitic(?) Smectite, ~60% →	60-40%. →	-do-, ± 75%.
Average Grain Size:	± 1 mm →	± 2mm →	~ 5 mm
Average Grain Shape:	angular →	-do- →	-do-
Sorting:	moderately poorly-sorted →	moderately-sorted →	poorly-sorted
Porosity:	Moderately low →	Low →	Moderately low
Permeability:	very low →	very low. →	very low
Alteration:	Total (smectitization, oxidation, hydration, zeolitization) →	-do- (palagonitization) →	-do-
Remarks:	Smectitized hyaloclastic tephra developed on surfaces of AOB pillow lava — gradation from pillow lava into dark reddish-brown palagonite (glassy chilled rind on pillow) into fragmental hyaloclastites. Demonstrates origin of hyaloclastites quite nicely!; MnO_2 deposit (dendritic microbistromatal ~0.5 mm thick) observed on one of my samples.	-do-, gradation is obvious	Serves as "cement" for basalt like 5DK3 and other volcaniclastics described here as Matrix Cement for Volcanogenic Breccia 5DK13
		Serves as substrate for knobby, warty, one-generation, 5 mm thick MnO_2 deposit which displays dendritic growth both internally and as supertopography.	≤ 0.5 mm patinate MnO_2 coating (dendritic)

13 14 15

Rock Type: Volcanogenic Breccia → - do - → - do -

Color: Yellowish-orange (10YR 7/6) (mean color) ≠ (matrix) Reddish-brown (10R 3/6) (cement) Olive-brown (5Y 3/4) → Moderate Brown (5YR ?/4

Texture: clastic / pyroclastic → pyroclastic / clastic ≠ → clastic

Framework Grains: mostly subangular pebble of 5DK3 composition, other more differentiated species, ~35%. → basalt clasts AOB, related to 5DK3 (±30%, 4 mm, avge range: ±0.5 mm – 4 cm) ≠ Palagonite clasts AOB, related to 5DK3? ±20%, 4 mm avge range: ±1 mm – 1 cm ≠ Crystals (enclosed) augite (3 ± 5%) ≠ avge size: ±2 mm dia. Nodrotinite / phillipsite ±5 mm → AOB fragments (clasts) (related to 5DK1?)

Cement: See 5DK12 → foraminiferal chalk, ±5% phyllo... 1 cm

Average grain size: ±3 cm → - do - !

Average grain shape: subangular → angular → - do -

Sorting: poorly-sorted → very poorly-sorted → - do -

Porosity: } See 5DK12 → Low → Moderate

Permeability: } → Very low → Moderate

Alteration: - do - → ≠ intense (oxidation, smectitization, zeolitization) → - do -

Remarks: - do - → - None - → appears to be monolithologic (AOB) Slope debris, (talus), very loosely-cemented, with many crevice-form openings between the talus fragments. clasts of the breccia are highly stained w/ $Fe(OH)_3$? and the breccia serves as substrate for thick (2 mm – 8 cm) knobby, warty-textured to laminar-dendritic (cyclical) one generation crusts of Mn...

16 17 18

ock Type: Volcanogenic Breccia → do → do

lor: Pale greenish-yellow (10Y 9/2) | Very Pale orange (10YR 8/2) | Yellowish-gray (5Y 7/2)

exture: Clastic → do → do

ame work rains: 1.) AOB intraclasts (prob. related to 5DK 2) (9%/of AOB)
Age size: ~5mm - 4cm (ice rafted)
Abundance: ±10%.
2.) AOB hyaloclastite → do
Age size: ~5mm
Abundance: ~10%.
→ do | but ~25% AOB clasts → do

ment: Coccolith(?) chalk, ±50% | do | but ~65% chalk, ±10% volcanic ash(?) | ~50% chalk, ~25% tephra

atrix: AOB sand, ±5%.
AOB hyaloclastite sand ±25% (related to 5DK14)

age Grainsize: ±2mm (range: ±0.1mm - 4cm) → do → do

ge grain Shape: Subangular → do → do

t. ng: Very poorly-Sorted → do → do

rosity: Moderately high | ± Very high → do

rmeability: Low | ± Low → do

teration: Superficial (phosphorization, zeolitization) → do → smectitization

rks: cobble-to-boulder-size clasts dislodged from outcrop by dredge; apparently developed on Volcanogenic breccia 5DK 14 as several pieces are in contact with this breccia and have it adhering to them; both the matrix and fragments of the manganitic cement of 5DK 14 are incorporated into this breccia, the manganite lining the coccolith chalk cement of 5DK 16 grades; surface of this breccia cement is, in turn, covered thinly (1-5 mm) with coccolith chalk which is, in turn covered with a patina of $Fe(OH)_3$(?) & MnO_2

do | Similar to 5DK 16 but lacking the manganite and AOB hyaloclastite intraclasts; appears to be developed on (& includes material from) a more tephra-rich breccia (5DK 18); serves as substrate for ±5mm-thick one generation dendritic growth (internally), Knobby, warty MnO_2 crust w/ finely corrugate superstopography | None

	19	20	21
Rock Type:	Volcanogenic Breccia	Phosphorized Chalk	Coquina Limestone
Color:	Very pale yellowish-orange (10YR 9/6)	very pale orange (10YR 9/2)	very pale yellowish-orange (10YR 9/6)
Structure:	Clastic	granular	clastic
Framework Grains:	Andesite clasts 20–40 % (extremely variable) Shell debris, 40–60% Intrastices, ± 15 %	foraminiferal tests, ≤ 100%	–do– ± 25 %, ≤ 1 mm dia Shell fragments, ± 20 %, ± 3 mm lg pellets, ± 45 %, ± 2 mm dia avg algal mat fragments(?) ± 10 %, ± 5 mm andesite fragments, < 1 %, ± 2 mm dia
Cement:	Calcite, recrystallized ± 5 %	–do– & inhomogeneous apatite	Calcite (matrix material apparently held together by calc. dissolved from itself and then recrystallized
Average Grain Size:	≈ 0.5 cm (range: ± 1 mm – 3 cm)	≤ 0.20 mm	± 2 mm
Average Grain Shape:	subangular	round	
Sorting:	Very poorly-sorted	Very well-sorted	moderately poorly-sorted
Porosity:	Extremely high	varies between dense and porous varieties (≈ 40–95 %)	Extremely high
Permeability:	Very high	– do – , (≈ 10 – 100 %)	Extremely high
Alteration:	superficial (solution, recrystallization of calcite; deposition of Fe(OH)? by downward diffusion from overlying ferromanganese crust.	Superficial (phosphorization of calcite, penetration by remobilized Fe(OH)$_3$(?) & Mn(OOH))	superficial (solution, redeposition recrystallization of calc. in matrix components
Remarks:	Cobble-to-boulder-size clasts broken from outcrop by dredge, composed largely of lagoonal-type shallow water carbonate debris (shells, pellets, oolites, coprolites, algal mats(?), corals, bryozoans, sponges, etc.) and volcaniclastics; serves as substrate for ± 5 mm-thick one generation, dendritic (external morphology), knobby, warty (external morphology) ferromanganese crust with coarsely nodose supertopography; ≈ 30 % of the knobs display effects of current scouring and are polished; calcareous material is stratified, and one piece (5DK 19b) displays a ± 2 cm-thick bed of much denser, volcaniclastics-poorer (only ≈ 7 %) description	very angular pebble- to cobble-size clasts broken, apparently, at some time before being collected by dredging (few surfaces are without Fe, Mn deposits); most pieces are intensely penetrated with tubular bore-holes (worms, clams?); these serve as conduits for passage of seawater and most are lined with apatite, either as a thin varnish-like deposit, and subsequently, Fe(OH)$_3$(?) and Mn(OOH)(?); in section, phosphorization may be seen as an aureolic development peripheral to the walls of the conduits and on surfaces once exposed to seawater; two types of this deposit are recognized: (a) porous, friable, light-weight, and highly penetrated with tubes; (b) denser, less porous, recrystallized, more replaced by apatite, less penetrated by tubes; surfaces exposed to seawater as outcrop	Cobble-size subangular clasts, perhaps dislodged from unexposed outcrop by dredge (no Fe or Mn. deposits anywhere); facies stratification apparent – probably deposited in a lagoonal environment or on a submarine shallow terrace related to 5DK19, but cleaner, w/o the volcaniclastic material.

	22	23	24
		Biotite Monzonite ?	Biotite Diorite ?
Rock Type:	Foraminiferal Chalk		
Color:	White (N9)	Dark greenish-gray (5GY 3/1)	→ -do-
Texture: Framework:	granular foraminiferal tests ± 99% ± 0.10 mm dia.	granitic equigranular (Phenocrysts) - Absent -	Gabbroic granular Hornblende? (Totally altered to biotite ± 50%, 3 mm avge dia.
Matrix:	foro Manganese micronodules ≤ 1% ≤ 0.10 mm dia.	(Groundmass) Plagioclase ± 30% Biotite ? ± 30% Hornblende ? ± 10% Augite (?) ± 10% K-feldspar ± 20%	Plagioclase, ± 50%, ± 2 mm dia.
Cement:	Probably lacking (rock is very soft and friable)		
Average grainsize:	≤ 0.10 mm	± 2 mm	
Average grainshape:	round		
Sorting:	Very well-sorted		
Porosity:	Extremely high		
Permeability:	Extremely high		
Alteration:	lacking (phosphorization?) negligible	Slight (oxidation, smectitization (?))	Intense (Biotitization of amph.; kaolinization of feldspar?)
Remarks:	Pebble-to-cobble-size fragments dislodged by dredge from outcrop, very similar in aspect to the phosphorized chalks (5DK20) but much less indurated, much more permeable and porous; penetrated by boring, burrowing animals (worms? clams?) making the rock cavernous, like 5DK20 (generally to a smaller and less diversified extent) The tube walls which penetrate the rock are stained slightly w/ $Fe(OH)_3$ (?), coloring them brownish-orange.	Single large cobble, 14 cm x 11 cm x 8 cm, subangular, loosened from hyaloclastite 5DK8, probably; possibly represents a loosened fragment of wall-rock from within the volcano or a fragment (xenolith) derived from a differentiated magma chamber below the volcano. the evenness of the grainsize indicates formation in a stable environment, free from influxes of fresh magma from other sources probably erupted subaqueously along with the hyaloclastite as a xenolith.	2, large pebble-size xenoliths from. → -do-, except this one appears to have been affected by entry of K-rich fluids (biotite coronas on amphibole)