RIDGE 29

S Karin Ridge

AMPLE ID	Mn CHEM	Pt & Au	REE	Si Chem	◎P.Sect.	35mm Pho	Archive	Paleo					
CD29-1													
CD29-2	[illegible]	[illegible]	[illegible]	✓	✓		✓		[illegible]	Slab to [illegible]	Si [illegible]		
CD29-3								✓					
CD29-4													
CD29-5													
CD29-6													
CD29-7	[illegible]				✓								
CD29-8													
CD29-9													
CD29-10													
CD29-11	[illegible]												
CD29-12													

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD 29-1 Location: S. Kashin Ridge

Size: 53 x 60 x 12 cm Weight: 55 Kg

Mn crust thickness:

Total: min: 75mm max: 130mm ave: 90mm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: subdued botryoidal to ripply

Internal structure:

layered: 4+
laminated:
massive: 4
porous:
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Yale Basalt

Description: Red brown to yellow brown altered vesic. basalt (5x3cm - 5mm. diam.) with green altered Mn oxide matrix. Some veins in fractures and vesicles

Mineralogy (XRD):

CD 29-1

Analyses and subsamples:

analysis:	analyst:
Kollman monument shop cut in half	11/88

	Max	Min	Ave	Texture.
layer 1	33	24 mm	28 mm	large pores at bottom of layer and very porous grades up the massive some of the pores are filled with clay
layer 2	25 mm	15 mm	20 mm	massive in center and complete; on either edge it is porous at bottom of layer
layer 3	30 mm	12 mm	20 mm	[illegible]
layer 4	50 mm	10 mm	20 mm	dense very fractured layer



Layer 1

Layer 2

Layer 3

Layer 4

Thinest part

Thickest part

CD29-1

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD 29-2 Location: S

Size: [illegible] Weight: 55kg Broken out in 4 parts A-D for photo — one piece sliced into 3 slabs

Mn crust thickness:

Total: min: 100 max: 130 ave: 115

	min:	max:	ave:
Inner crust:		1 - 10 2A	ave: 15
Outer crust:		2B 25 3	ave: 30
		4	30

Surface texture: botryoidal w/ pelletoid [illegible]

Internal structure:

layered: 4; 1 w/ 2 sub layers
laminated: 4 dense, fractured
massive: 3 micro-fractured
porous: 2A, pillar structure
dendritic:
other: 2B Dense, pillar structure
1 crudely laminated w/ rock frag. inclusions

Mineralogy (XRD): 1, 2A, +3 have Fe-oxide staining

Associated alteration, phosphorite, or hydrothermal deposits: + fracture fill
Brown phosphorite (Burrows Hill?) in host Rx — also vein + Fracture fill in Crust layer 4

Substratum:

Rock type: Volc. Breccia - hyaloclastite

Description: see CD29-1
pale yellowish-brown - w/ phosphate filling veins & replacing carbonate layers

Carbonate rock mixed in w/ hyaloclastite

Mineralogy (XRD): Three small slabs cut for PTS if no one else wants them

Layers 1+2 and 3+3 split easily along vertical discontinuities — layers 3+4 [illegible]



CD29-2

Analyses and subsamples:

analysis:	analyst:
Kollman monument shape cut in half	1/6/88
CD29-2A Bulk (CC) PTS Pt, Au	0-110 mm 110 mm (105g)
CD29-2B outer CC, Pt, Au REE	0-9 mm 9 mm (6g REE) (43g rest)
CD29-2C - layer 2A+B CC, Pt Au, REE	9-47 mm 40 mm (21g REE) (102g rest)
CD29-2D - layer 3 CC, Pt Au, REE	47-76 mm 27 mm (10g REE) (91g rest)
CD29-2E - Bulk (31g) CC	Substrate - hyal-phos
Various Layers Sr, Pb isotopes	Hein
Piece coated for display	
1 Slab, Be, Pb, Sr, Nd, U isotopes	Hein, O'Nions
Slab 5 sent to Lalocui	
2 other slabs archived	
CD29-2A Pt, Ir - Hz	
CD29-2E - layer 4 CC, Pt Au, REE	75-105 mm 30 mm (15g REE) (90g rest)
1 Slab - Godfrey - Hf isotopes	
1 Slab - Chan? Li isotopes	Slab to Gallic

CD29-2

CD29-2 All pieces from #2

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: Sta 31 CD 29-3 Location: S Kerin Ridge

Size: 38 x 19 x 13 cm Weight: 35 Kg

Mn crust thickness:

Total: min: 80 mm max: 120 mm ave: 100 mm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Subdued botryoidal ridged; granular [illegible] bottom. Many burrows in [illegible]

Internal structure:

layered: 5 ?
laminated: uppermost, 3rd & 4th
massive:
porous: 2nd inner most
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

minor phosphate veining in inner most Mn layer

Substratum:

Rock type: Volc Breccia

Description: Orange-red angular altered pelagic(?) clasts 1 mm² - 1 cm² in Mn-rich altered [illegible] matrix

fine grained [illegible]

Phosphorite clasts (?) in upper 2nd to last crust

1 layer (or more) red [illegible] Foram sand between crusts [illegible] Recent? and maybe a [illegible]

Mineralogy (XRD):



CD29-3

Analyses and subsamples:

analysis:	analyst:
Paleo on Foram sand	

	Max	Min	Ave	
Layer 1	10	8 mm	10 mm	Massive
Layer 2	22.	15 mm	20 mm	large [illegible] at bottom of layer, porous throughout the layer
Layer 3	40 mm	30 mm	25 mm	Porous, fractured, denser and [illegible] top of layer.
Layer 4	43 mm	30 mm	35 mm	massive at base of layer; fractured, growth lamellae [illegible] Sample

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: Stn 31 CD 29-4 Location: S Karin Ridge

Size: 41x31x15cm Weight: 23 Kg

Mn crust thickness:

Total: min: 110mm max: 140mm ave: 128mm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Subdued botryoidal w/protuberances of smaller botryoids, granular

Internal structure:

layered: 4(+)
laminated: upper, 2nd, inner most
massive:
porous: 3rd
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Volc Breccia

Description: Yellow orange stained clasts in yellow Mn-rich matrix. Almost no substrate - rocky just a surface coat

Mineralogy (XRD):

CD 29-4

Analyses and subsamples:

analysis:	analyst:

	Max	Min	Ave	
Layer 1	12mm			massive
Layer 2	45mm	40mm	45mm	porous, [illegible] separating sublayers, above is more porous [illegible]
Layer 3	25mm	15mm	18mm	[illegible]
Layer 4	70mm	30mm	[illegible]	Massive [illegible] Fractures growth lamina [illegible]

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: Sta 31 CD 29-5 Location: S Kaena Ridge

Size: [illegible] 15 cm Weight: 40 Kg

Mn crust thickness:

Total: min: 9 max: 16 ave: 12.5 cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Sediment botryoidal w/nodules

Internal structure:

layered:
laminated:
massive: ✓
porous:
dendritic:
other:

CD29-5

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: None

Description:

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:

	Max	Min	Ave	Texture
Layer 1	55	50mm	53mm	Porous, vesicular surface mineralogy [illegible] Rough large pores (Not Filled) in Rough [illegible] middle of layer
Layer 2	35mm	30mm	33mm	Fe Stained
layer 3	50mm	30mm		Dense Massive, Fractured with [illegible] terminus visible

Layer one = layer from [illegible]

CD29-5

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31-CD29-6 Location: S. Kane Fracture Zone

Size: 31x29x6 Weight: 2.5 kg

Mn crust thickness:

Total: min: ____ max: 115 ave: ____

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: Ripples to massive, botryoidal

Internal structure: ____

layered: ____
laminated: ____
massive: ✓
porous: ✓ to a few mm
dendritic: ____
other: ____

Mineralogy (XRD): ____

Associated alteration, phosphorite, or hydrothermal deposits:

phosphorite crust

Substratum: Very little

Rock type: ____

Description: ____

Mineralogy (XRD): ____



CD29-6

CD29-4 Phos Study

Analyses and subsamples:

analysis:	analyst:
38mm Slide (missing Phos bleb)	Sm Ah

	Max	Min	Ave	Texture
Layer 1	10mm	[illegible]	[illegible]	[illegible], laminated
Layer 2	25mm	20mm	[illegible]	Dendritic w/ large pores at bottom of layer (clay filled)
Layer 3	25mm	20mm	23mm	pillar structures, laminated mass [illegible] at the bottom of the layer a few larger clay filled pores [illegible]
Layer 4	28mm	25mm	[illegible]	Fe stained, dendritic
Layer 5	35mm	20mm	33mm	Massive, Porous, Fractured layer; laminated

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31-CD29-7 Location:

Size: 31x20x16 cm Weight: 10 kg

Mn crust thickness:

Total: min: 110 mm max: 150 mm ave: 130 mm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Smooth to botryoidal (nodular)

Internal structure:

layered: 4
laminated:
massive: innermost, outermost
porous: 2 & 3
dendritic: 4th
other: 2 is pillar structure

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Phosphatic veins in crust 1 & 2 to Mn crust - mostly in innermost massive-Dense layer

Substratum: very little

Rock type: Vole Breccia - hyaloclastite

Description: Yellowish hyaloclastite

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:
CD29-7A (cc) 0-80	80 mm (layers 1-4 (103 g)
CD29-7B (cc) 80-138	58 mm (layer 5 (48 g)
layer 5 - Dense, vitreous	Subconcoidal fracture
DPTS 0 - 130 mm (3 or 4 CD29-7C	sections)
layer 3 has recent mud in it - most removed	

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD29-8 Location: [illegible]

Size: 31x18x14cm Weight: ≈ 10 Kg

Mn crust thickness:

Total: min: 80mm max: 90mm ave: 80 mm

Inner crust: min: ___ max: ___ ave: 3.5cm

Outer crust: min: ___ max: ___ ave: 6 cm

Surface texture: Slightly botryoidal

Internal structure: ___

layered: ___
laminated: 2 cm
massive: ___
porous: ✓ layer 2
dendritic: ___
other: ___

Mineralogy (XRD): ___

CD29-8

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Volc. Breccia

Description: Yes, altered vesicular and [illegible] along fracture

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:

	Max	Min	Ave	
Layer 1				
Layer 2	35mm	20mm	30mm	Porous
Layer 3	15mm	15mm	15mm	Fe-Stained
Layer 4	35mm	25mm	30mm	Dense, Massive, Fractured

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD 29-9 Location:

Size: Weight: 3kg nodules

Mn crust thickness:

Total: min: 2mm max: 15mm ave: 8mm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: perfectly smooth

Internal structure:

layered:
laminated:
massive: ✓
porous:
dendritic:
other:



Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Phosphatized Volcanic Breccia w/Mn crust

Description: Red brown to yellow phosphatized volcanic clasts with cream colored phosphate cement (upper 2 cm) and beige-brown calcite cement (lower 2 cm). Fragments of rock are Mn crust as follows: 30 mm thick, 2 layers: upper - dendritic, porous, botryoidal; lower - massive w/phosphatized vein overgrowths

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:



CD 29-9

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD 29-10 Location: S Kodiak Ridge

Size: _____ Weight: 8 kg

Mn crust thickness:

Total: min: 6cm max: 9cm ave: 7.5 cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: Subdued Botryoidal

Internal structure: _____

layered: 5
laminated: _____
massive: 3
porous: _____
dendritic: 1
other: _____

CD29-10

Mineralogy (XRD): _____

Associated alteration, phosphorite, or hydrothermal deposits: _____

Substratum:

Rock type: Nodule

Description: _____

Mineralogy (XRD): _____

Analyses and subsamples:

analysis:	analyst:
half given to Hubach	

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD 29-11 Location: S Kasei Valles

Size: 24 x 30 x 8 cm Weight: 4 kg

Mn crust thickness:

Total: min: 30 mm max: 40 mm ave: 35 mm

Inner crust: min: — max: — ave: —

Outer crust: min: — max: — ave: —

Surface texture: perfectly smooth, irregular in recesses.

Internal structure: Massive + highly fractured in

layered: areas, highly porous in other areas -

laminated:

massive: 1

porous:

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

phospho- Cement + veins, Mn-oxide Cement + fragments in the phosphorite -

Substratum:

Rock type: Porcellanite basalt breccia

Description: Very porous (vesicular) (1mm² - 1x3 cm) angular (lithic fragments) loosely cemented in phosphatic ~~Carbonate~~ mud. Clasts are primarily vesicular basalt (minor) and ~~a~~ few ~~lithic~~ lithic clasts in phospho- -

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:
CD 29-11, Bulk (CC)	0-35 mm, 35 mm
one layer has lot recent mud, may not have gotten it all ?	

Sample Description

Cruise I.D: F7-86-HW Sample I.D: Sta: 31 CD-29-12 Location:

Size: 19x15x12cm Weight: 4kg

Mn crust thickness:

Total: min: 4mm max: mm ave:

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: patchy dusting

Internal structure:

layered:
laminated:
massive:
porous:
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Volcanic Breccia

Description: Red brown altered volcanic fragments 75-½ mm [illegible] (small fragments [illegible] and rims of some [illegible] with phenocrysts [illegible] Clasts: Breccia is [illegible] groundmass [illegible]

Mineralogy (XRD):

CD29-12

Analyses and subsamples:

analysis:	analyst:

Cruise Id: EW-86-HW Size _____ (cm) Location: Karin Ridge

Sample Id: CD29-12 Weight: _____ (kg) Depth: _____ (m)

Mn Crust Description

Surface Texture: _____

Mn Crust Thickness (mm):

Min: _____ Max: _____ Ave: _____

Layers (Outer to Inner)(mm):

	Min	Max	Ave	Texture & Color
1.				
2.				
3.				
4.				
5.				
6.				

* Alicé has this sample. She has numbered it CD29-A4

Comments and XRD Mineralogy:

ANALYSES

Subsample Number	Analysis	LayerThickness

Substrate Description

Rock Type: Basalt (Alkalic)

Description:

Biology ?: _____

Described By: Alicé Davis

Subsampled By: