Dredge 7

L5-83 HW
D7

D7-A1

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D 7-A1

Location: Necker Ridge

Size: 6 x 5 x 3 cm Weight:

Mn crust thickness:

Total: min: max: ave: 1.5 cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture:

Internal structure:

layered:
laminated:
massive:
porous:
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: ?

Description:

Mineralogy (XRD):

See picture for piece

May 1998: crumbly, highly fractured 5 x 7 x 3.5 cm piece left, w/ several small, broken pcs ≈ 1 x 2 x 2 cm each

Analyses and subsamples:

analysis:	analyst:
1 piece 2½ x 3 cm thick to:	Bob Willach Minneapolis, Minn
1 piece to: for flotation separation	Don. G. Foot Jr. Salt Lake City, Utah
1 piece	Lynn Ingram
Pt group (12/86)	Floyd Bonnar
Se Won Chang took subsample 10	
May 1998 sent pc. to Alex Halliday	

Uniform pr → skew results
Chem – Bad past 53 cut to match PK
lg PK = larger atom –
sm PK = smaller atom

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7-A2

Location: Necker Ridge

Size: 50x32x17cm Weight: 19Kg

Mn crust thickness:

Total: min: 3.5 max: 4cm ave: 3.8cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: smooth nodular botryoidal

Internal structure:

layered:
laminated: ✓
massive:
porous: ✓
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Vole. Breccia

Description: Hard altered red-brown vole. clasts, sandy matrix (1mm-5cm); basalt is red to brown altered yellow sandy matrix

Mineralogy (XRD):

L5-83-HW

D7-A2

Analyses and subsamples:

analysis:	analyst: (or sent to sampled by)
end piece given to. 19x17x15cm	John Toth Analytical Services
piece to	A. Davis
REE (Denver) and CC (Boston) of MW crusts	Jung-Keuk Kang 6/8
I 0-3 mm 4gm (CC)	Job # SW32 (REE
II 3-8 " 3 "	" # BZ89 (CC
III 8-16 " 6 "	
IV 16-26 " 5 "	
V 26-36 " 2 "	

Archive Splits

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D 7-A3 Location: Necker Ridge

Size: 36x36x8 cm Weight: _____

Mn crust thickness:

Total: min: ____ max: ____ ave: 3.8cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: smooth - slight botryoids

Internal structure: _____

layered: ✓

laminated: _____

massive: _____

porous: _____

dendritic: _____

other: _____

Mineralogy (XRD): _____

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Volc. breccia

Description: Highly altered red-orange, sub-rounded, poorly sorted (1mm-2cm) lithic volc. clasts in an altered yellow, zeolitized tuffaceous sand matrix

Mineralogy (XRD):

Cut & broken up - pieces missing

L5-83-HW
D7-A3

Analyses and subsamples:

analysis:	analyst:
XRD:	C. Daniel
I. Substrate-composite	
II outer crust	
III inner crust	
Doubly Polished Thin Section	
Chemistry-substrate	
Density	

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7-A5

Location: Necker Ridge

Size: [illegible] Weight: 2462g

Mn crust thickness:

Total: min: 1.5 max: 3cm ave: 2cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: Botryoidal, fibrous, smooth

Internal structure: ____

layered: ____
laminated: ____
massive: ____
porous: ____
dendritic: ____
other: ____

Mineralogy (XRD): ____

Associated alteration, phosphorite, or hydrothermal deposits:

Fe oxides in Mn crust

Substratum:

Rock type: Mafic Breccia

Description: Highly altered, yellow to brown, fractured manganese cemented, poorly sorted (1-10cm), tightly packed, volcanic clasts in a yellow, poorly sorted silty sand matrix, w/ pockets of tan, silty sediment, white, calcareous sediment.

Mineralogy (XRD):

D7-A5 L5-83-HW

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-1 Location: Decker Ridge

Size: 15x10x5cm Weight: ~3000g

Mn crust thickness:

Total: min: 3 max: 15 ave: 4cm

Inner crust: min: 2 max: 2.5 ave: 2cm

Outer crust: min: 1 max: 1.5 ave: 1cm

Surface texture: Smooth, flat

Internal structure:

layered: ✓ ~4

laminated:

massive:

porous:

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Red crust is overlain with a thin layer of sediment

Substratum: NONE

Rock type:

Description:

Mineralogy (XRD):

LS-83-HW

D7-A8-1

Analyses and subsamples:

analysis:	analyst:
XRD:	Rosemary ?
inner	
outer	

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-2 Location: Necker Ridge

Size: [illegible] Weight: [illegible]

Mn crust thickness:

Total: min: 3cm max: 3.5cm ave: 3cm

Inner crust: min: 2.4cm max: 2.8cm ave: 2.6cm

Outer crust: min: 6mm max: 8mm ave: 7mm

Surface texture: [illegible]

Internal structure:

layered: ✓
laminated:
massive:
porous: ✓
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Fe oxide in crust

Substratum: [illegible]

Rock type: [illegible]

Description: Altered yellow-brown basaltic sub rounded pebble (~4cm) highly packed volcanic clasts in a hard (silicified? to incipient?) sandy matrix with white calcareous sediment on bottom

Mineralogy (XRD):

L5-83-HW
D7-A8-2

Analyses and subsamples:

analysis:	analyst:
Pt Core (1/4D)	Floyd Rowe
D7-A8-2A	
D7-A8-2B	

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7-48-3

Location: Necker Ridge

Size: 20x 2x6cm Weight: 2 kg

Mn crust thickness:

Total: min: 3.0 max: 3.5 ave: 3.5cm

Inner crust: min: ___ max: ___ ave: 2.7cm

Outer crust: min: ___ max: ___ ave: 2mm

Surface texture: Smooth, rounded

Internal structure:

layered: ✓ ?

laminated:

massive: ✓

porous: ✓

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Fe oxides in crust

Substratum:

Rock type: Volc. breccia?

Description: Altered ? rounded pebble? breccia? cemented with ferric? clasts ? basalt with ? equigranular matrix. Thin ? phosphate? layer between Mn crust & substratum

Mineralogy (XRD):

L5-83-HW

D7-48-3

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7 48-4 Location: Necker Ridge

Size: small hcm Weight: _____

Mn crust thickness:

Total: min: 2.5 max: 4 ave: 3.5cm

Inner crust: min: _____ max: _____ ave: _____

Outer crust: min: _____ max: _____ ave: _____

Surface texture: Porous & smooth

Internal structure: _____

layered: _____
laminated: _____
massive: _____
porous: ✓ extremely
dendritic: _____
other: _____

Mineralogy (XRD): _____

Associated alteration, phosphorite, or hydrothermal deposits:

Ferruginous crust

Substratum: thin layer

Rock type: talc breccia

Description: Altered red to brown SLC rounded, poorly sorted (1-4cm) loosely packed volcanic clasts in a yellow-brown, well indurated sandy matrix speckled w/ Mn & small pockets of white phosphatic material

Mineralogy (XRD): _____

L5-83-HW
D7-48-4

Analyses and subsamples:

analysis:	analyst:
XRD	C. Daniel
D7-48-4-I sf. crust	
II 1.5-5.0 cm	
Doubly polished thin section	

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D 7-A8-5

Location: Necker

Size: 20x8x10cm Weight: 2.03 kg

missing! 9-24-85

Mn crust thickness:

Total: min: max: ave:

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture:

Internal structure:

layered:
laminated:
massive:
porous:
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type:

Description:

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:
2/3rx given to:	Buford Hall MMS 1 Golden Shore Suite 260 Long Beach, Ca. 90802

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-6 Location: Necker Ridge

Size: 21x16x6cm Weight: ~2.5kg

Mn crust thickness:

Total: min: ____ max: ____ ave: 3cm

Inner crust: min: ____ max: ____ ave: 2.5cm

Outer crust: min: ____ max: ____ ave: 0.5cm

Surface texture: Smooth

Internal structure: ____

layered: ✓ 2

laminated:

massive:

porous:

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum: thin layer

Rock type: Volc. Breccia?

Description: Sub-rounded volcanic clasts coated w/ Mn cemented w/ phosphate

Mineralogy (XRD):

L5-83-HW

D7-A8-6

Analyses and subsamples:

analysis:	analyst:
XRD	C. Daniel
D7-A8-6-I bulk crust	
6-II bulk substrate	

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7 A8-7 Location: Necker Ridge

Size: 9X12X8cm Weight: ≈500g

Mn crust thickness:

Total: min: ____ max: ____ ave: 3.5cm

Inner crust: min: ____ max: ____ ave: 3cm

Outer crust: min: ____ max: ____ ave: 0.5cm

Surface texture: Smooth

Internal structure: ____

layered: ✓ in 4

laminated:

massive:

porous: ✓

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Fe Oxide Staining

Substratum: thin layer

Rock type: Volc. Breccia

Description: One large (>6cm round) brown volc. clast with a few smaller ones loosely packed in a yellow, zeolitized, silty sand matrix

Mineralogy (XRD):

L5-83-HW

D7-A8-7

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7 A8-8

Location: Nectar Ridge

Size: 11x6x6cm Weight: ~250 g

Mn crust thickness:

Total: min: 3cm max: 5cm ave: 3.5cm

Inner crust: min: ____ max: ____ ave: 2cm

Outer crust: min: ____ max: ____ ave: 1.7cm

Surface texture: Smooth to porous

Internal structure: ____

layered: ✓ ~2-3

laminated:

massive: ✓

porous: ✓ columnar

dendritic:

other:

L5-83-HW

D7-A8-8

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum: very little

Rock type: Volc. breccia

Description: a few volc. clasts coated w/Mn & cemented w/an orange-brown silty sand sediment

Mineralogy (XRD):

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-9 Location: Nector Ridge

2 pieces: 7x8x5cm

Size: 9x7x5cm Weight: ~500g

Same as D7-A8-4

Mn crust thickness:

Total: min: ____ max: ____ ave: 3.5cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: extremely porous

Internal structure: ____

layered: ____
laminated: ____
massive: ____
porous: / extremely
dendritic: ____
other: ____

Mineralogy (XRD): ____

Associated alteration, phosphorite, or hydrothermal deposits: ____

Substratum: same as D7-A8-4

Rock type: ____

Description: ____

Mineralogy (XRD): ____

L5-83-HW
D7-A8-9

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A3-10 Location: Decker Ridge

Size: _____ Weight: ~1 kg

Mn crust thickness:

Total: min: ____ max: ____ ave: 3cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: _____

Internal structure: _____

layered: _____
laminated: _____
massive: _____
porous: _____
dendritic: _____
other: _____

Mineralogy (XRD): _____

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: _____

Description: _____

Mineralogy (XRD):

Bag of small pieces of crust w/no substratum

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7A8-11 Location: Decker Ridge

Size: _____ Weight: 2.19

Mn crust thickness:

Total: min: ____ max: ____ ave: 3cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: Smooth grainy botryoidal

Internal structure: _____

layered: _____
laminated: _____
massive: _____
porous: _____
dendritic: _____
other: _____

Mineralogy (XRD): _____

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: rock breccia

Description: _____

Mineralogy (XRD):

Bag of small broken pieces of Mn crust w/ very little or no substratum

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-12 Location: Juan de Fuca Ridge

Size: 8x4cm Weight:

13.7x5cm

Mn crust thickness:

Total: min: max: ave: 1mm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture:

Internal structure:

layered: ✓ ?

laminated:

massive:

porous: ✓

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Vitric breccia

Sample w/ thin crust has very little substratum

Description: Altered, lithified, subrounded, poorly packed volcanic clasts (~1cm) in a yellow-orange cement and some mn crust

Mineralogy (XRD):

L5-83-HW

D7-A8-12

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-13 Location: Vents Ridge

Size: 5 small pieces Weight: 2 kg

Mn crust thickness:

Total: min: ____ max: ____ ave: 2cm

Inner crust: min: ____ max: ____ ave: ____

Outer crust: min: ____ max: ____ ave: ____

Surface texture: Smooth & porous

Internal structure: ____

layered: ____
laminated: ____
massive: ____
porous: ____
dendritic: ____
other: ____

Mineralogy (XRD): ____

L5-83-HW
D7-A8-13

Associated alteration, phosphorite, or hydrothermal deposits: ____

Substratum:

Rock type: Vole Breccia
2 have no substratum

Description: Breccia, vole clast (8x5cm) w/ porous Mn oxide w/ red orange Fe oxide coating. Other 2 rocks have a white phosphate layer between crust & breccia w/ extensive red orange Fe oxide staining on breccia.

Mineralogy (XRD): ____

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7-A8-14

Location: Necker Ridge

Size: Harrison Weight: 25000g

Mn crust thickness:

Total: min: max: 4.5 ave: 3cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Smooth & porous

Internal structure:

layered: ✓

laminated:

massive: ✓

porous: ✓

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum: very little

Rock type: Vol. Breccia w/phosphate

Description:

Mineralogy (XRD):

L5-83-HW

D7-A8-14

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW

Sample I.D: Sta: D7-A8-15 Location: Necker Ridge

Size: 7pieces 18x10x5cm (largest) Weight: 2.2 kg

Mn crust thickness:

Total: min: max: ave: 3cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: rough, porous, botryoidal

Internal structure:

layered: ✓ ~2

laminated:

massive: ✓

porous: ✓

dendritic:

other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum: (not much substratum)

Rock type: Volc breccia

Description: Highly altered yellow hyaloclastite? 2 to a few small (1-2cm) volc. clasts. The largest sample has thin layer of phosphate

Mineralogy (XRD):

L5-83-HW

D7-A8-15

Analyses and subsamples:

analysis:	analyst:
XRD:	C. Daniel
I: substratum	
II: porous crust (0-3cm)	
Chemistry-substratum	
Chemistry-bulk (of porous crust)	
Pt group (12/86)	Floyd Brown-Boston
Au analysis (12/86)	Dave McKown-Denver

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D7-A8-16 Location: Nazca Ridge

Size: 5 pieces 4x8x4cm (largest) Weight: 2.5kg

Mn crust thickness:

Total: min: ___ max: ___ ave: 2cm

Inner crust: min: ___ max: ___ ave: ___

Outer crust: min: ___ max: ___ ave: ___

Surface texture: Smooth & Flat

Internal structure: ___

layered: ___
laminated: ✓
massive: ___
porous: ___
dendritic: ___
other: ___

L5-83-HW
D7-A8-16

Mineralogy (XRD): ___

Associated alteration, phosphorite, or hydrothermal deposits: ___

Substratum:

Rock type: Lithified Talus Breccia

Description: Brown altered subrounded volcanic clasts (1-3cm) in a matrix of tan phosphate cement

Mineralogy (XRD): ___

Analyses and subsamples:

analysis:	analyst:

Sample Description

Cruise I.D: L5-83-HW Sample I.D: Sta: D 7-B1 Location: Necker Ridge

* Size: 20 cm across Weight: ~500g
Already cut & piece taken

Mn crust thickness:

Total: min: 1.0 max: 2.3 ave: 1.8cm

Inner crust: min: max: ave:

Outer crust: min: max: ave:

Surface texture: Botryoidal across

Internal structure:

layered:
laminated:
massive:
porous: ✓
dendritic:
other:

Mineralogy (XRD):

Associated alteration, phosphorite, or hydrothermal deposits:

Substratum:

Rock type: Basalt to Vole Breccia

Description: One half of rock is a gray-brown basalt clast, strongly altered (percentage of olivine?) other half is an altered yellow to orange cemented volcanic breccia with well sorted (~3cm) clasts packed & fractured in a yellow (zeolitized) silty matrix

Mineralogy (XRD):

LS-83-HW
D7-B1

Analyses and subsamples:

analysis:	analyst:
XRD	C Daniel
D7-B1-I bulk crust	
D7-B1-II bulk substrate	
Doubly Polished Thin Sec	Mann petrographic