

STRATIGRAPHIC NOMENCLATURE—LOS ANGELES SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California.</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>Formally named formations grouped in sequence (separated by semicolons) are listed from youngest to oldest.</small>	
QUATERNARY	Qs	RECENT DUNE SAND	Wind-blown sand and beach deposits along coast, desert dune sand and bars of wave-deposited sand around Rosamond Dry Lake.	
	Ql	QUATERNARY LAKE DEPOSITS	Playa clay and silt of Rosamond Dry Lake.	
	Qal	RECENT ALLUVIUM	"Younger" alluvium consisting of Recent clay, silt, sand and gravel, unconsolidated, poorly stratified to well stratified, includes alluvial fan, flood-plain, and streambed deposits. In some desert areas includes mixture of playa clay and wind-blown sand.	
	Qf	RECENT ALLUVIAL FAN DEPOSITS IN THE GREAT VALLEY	Alluvial fan deposits at the southern end of the San Joaquin Valley.	
PLEISTOCENE	Qt	QUATERNARY NONMARINE TERRACE DEPOSITS	Alluvial terrace deposits, in places along the coast overlies thin marine terrace deposits. Locally includes fanglomerate and "older" alluvial debris at higher levels.	
	Qc	PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS	Castas Formation—red to greenish clay, silt, sand and conglomerate (Carpinteria area). Frazier Mountain Formation—conglomerate, fanglomerate and lamella breccia of local derivation (Lockwood Valley); Pacoima Formation—dark brown breccia and fanglomerate, locally tuffaceous; San Gabriel Formation—red to greenish clay, silt, sand and conglomerate (Pearland quadrangle). Harold Formation—interbedded gravel, sand, silt, and gyttiferous clay. "Older" alluvium consisting of gravel, sand, clay, and silt; dissected alluvial fan deposits. "Ancient" alluvium, commonly with soil formation on surface (San Gabriel Valley).	
	Qm	PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS	San Pedro Formation—sand, silt, and marl, highly fossiliferous (Los Angeles basin). Marine and some stream terrace deposits along the coast and on Santa Cruz Island. Qm-Pu: Santa Barbara Formation—buff to yellow sand, silt and calcareous sediments.	
	Qp	PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS	Paso Robles Formation—interbedded gray to brown clay, sandstone, and conglomerate (Cuyama area and Santa Ynez Mountains). Saugus Formation—sandstone, conglomerate and siltstone, mostly fuculite but locally brackish water deposits. Tulare Formation—gravel, clay and conglomerate (south end San Joaquin Valley).	
	Puc	UNDIVIDED UPPER PLIOCENE NONMARINE SEDIMENTARY ROCKS	Hungry Valley Formation—brown conglomerate, white conglomeratic sandstone, and brown mudstone.	
	Pc	UNDIVIDED PLIOCENE NONMARINE SEDIMENTARY ROCKS	Morales Formation—gray, poorly bedded stream-laid gravel and sand; Quatal Formation—orange to buff, pebbly, arkosic sandstone and light-red clay, gyttiferous siltstone, and gypsum; Lockwood Clay of Carman, 1964—pale-tan to gray montmorillonitic clay and interbedded sandstone, Peace Valley Formation—brown sandstone, buff sandstone, some conglomerate and sedimentary breccia. Meek Mine Formation—pebble gravel, sand, and lacustrine clay shale; may be early Pleistocene in age (Tehachapi Mountains area).	
	Pu	UPPER PLIOCENE MARINE SEDIMENTARY ROCKS	Sunshine Ranch Member of Saugus Formation—continental and brackish-water greenish sandstone and mudstone, red conglomerate beds, and thin limestone beds (San Fernando Reservoir area); upper member of the Fernando Formation (includes rocks commonly called Pico Formation)—sandy siltstone, conglomerate and sandstone (Los Angeles basin); Pico Formation—siltstone, sandstone and conglomerate (Ventura basin). Careaga Sandstone—poorly consolidated, fine- to medium-grained sandstone (Santa Ynez Mts.). Unnamed marine Pliocene-Pleistocene (?) silt and sand in the Baldwin Hills area.	
	Pmlc	MIDDLE AND/OR LOWER PLIOCENE NONMARINE SEDIMENTARY ROCKS	Anaverde Formation—gray shale, arkosic sandstone, red conglomerate, dark-gray dioritic breccia; plant fossils of early to middle Pliocene age (in San Andreas fault zone). Ridge Route Formation—sandstone with conglomerate, shale, and siltstone interbeds. Chanac Formation—sandstone, claystone and conglomerate.	
	Pml	MIDDLE AND/OR LOWER PLIOCENE MARINE SEDIMENTARY ROCKS	Lower member of the Fernando Formation (includes rocks commonly called Repetto Formation)—siltstone, conglomerate and fine sandstone. Echequin Formation—buff, fine, fossiliferous sand (SW San Joaquin Valley). Unnamed siliceous mudstone in the Bitterwater Creek area (may be late Miocene in age). Towsley Formation—siltstone, mudstone, sandstone and conglomerate (late Miocene and early Pliocene age; Santa Susana Mountains).	
	Mc	UNDIVIDED MIOCENE NONMARINE SEDIMENTARY ROCKS	Fiss Fanglomerate—fanglomerates of brown volcanic detritus and gray granitic detritus (Miocene age uncertain; Willow Springs and Rosamond quadrangles). Punchbowl Formation—white, buff to pink sandstone, gray to red siltstone and clay shale, and gray to red conglomerate (middle and late Pliocene age; Valley area); Yuba Formation—Mint Canyon Formation—grayish-green to light-brown siltstone and mudstone, light-brown, gray or reddish sandstone and conglomerate; some tuff beds (late Miocene and early Pliocene age; Mint Canyon area). Oso Canyon Formation—fanglomerate, gray-white to red sandstone and red green siltstone, grades laterally west into Quail Lake Formation (late Miocene age; west end Antelope Valley). Bissell Formation—dolomite, limestone and shale with magnetite layers, sandstones and conglomerates (Miocene or Pliocene age; Bissell Hills, Rosamond quadrangle). Unnamed gray fanglomerate sand and clay that grades northward into marine sandstone and shale of middle and late Miocene age (southern San Joaquin Valley).	
Mu	UPPER MIOCENE MARINE SEDIMENTARY ROCKS	Tequepis Sandstone—gray-white, massive to thick-bedded, compact, semitransparent sandstone (Santa Ynez Mts.). Sisquoc Formation—soft, fissile to massive dolomite in part tuffaceous shale and siltstone (Santa Ynez Mts.). Santa Margarita Formation—sandstone, siltstone, some conglomerate (Cuyama Valley and Pine Mtn. areas). Modelo Formation—siliceous and diatomaceous shale, sandstone, siltstone and some conglomerate (in part middle Miocene; Santa Monica Mts.). Upper part of Monterey Shale on Pt. Dume, Castaic Formation—shale with interbedded sandstone and beds of pebble conglomerate (considered to be same as the Modelo Formation by some geologists). Puente Formation—siltstone, siliceous shale, diatomaceous shale, sandstone and conglomerate, including Sycamore Canyon Member—conglomerate and siltstone (Los Angeles area). Quail Lake Formation—buff sandstone and brown shale (west end Antelope Valley).		
Mmc	MIDDLE MIOCENE NONMARINE SEDIMENTARY ROCKS	Tick Canyon Formation—siltstone, sandstone, conglomerate, probably early Miocene. Unnamed nonmarine sandstone, mudstone and conglomerate and Tick Canyon Formation of Durrell, 1954, Santa Monica Mts.). Unnamed cobble conglomerate and red beds (southern San Joaquin Valley).		
Mm	MIDDLE MIOCENE MARINE SEDIMENTARY ROCKS	Monterey Shale—brown, soft, fissile, punky, organic shale, hard porcellanous shale and chert, bituminous silty siliceous shale, locally includes thin limestone beds, siliceous clay siltstone and sandstone; south of Malibu Coast fault includes shale, sandstone, dolomite, and chert. Unnamed sandstone, mudstone, sedimentary breccia and some volcanic rocks (south of Malibu Coast fault). San Onofre Breccia—sandstone and breccia of schist fragments including stabs and blocks of glauconitic schist (along coast, western Santa Monica Mts.). Topanga Formation—shale, sandstone, some conglomerate, arkose (Santa Monica Mts.).		
TERTIARY	MIOCENE VOLCANIC ROCKS:			
	Mv	UNDIFFERENTIATED	Interbedded agglomerate, flow breccias, flows, tuffs, and volcanic sandstone (central Santa Monica Mts.). Andesite and basalt flows in the Topanga Formation(?), Verdugo Mts. and Pacoima Hills.	
	Mv ^a	ANDESITIC	Conejo Volcanics—hypersthene andesite (Santa Monica Mts.).	
	Mv ^b	BASALTIC	Conejo Volcanics—basalt flows and breccia, some andesite, arkose and tuff (Santa Monica Mts.). Vesicular and porphyritic extrusive basaltic flows and pyroclastic rocks (Conejo Volcanics); Santa Cruz and Anacapa Islands; Basalt flows in the Gem Hill Formation (Miocene age uncertain; Rosamond quadrangle). Vesicular amygdaloidal basalt flows, with interbedded tuff breccia and agglomerate (Sunland area).	
	Mvp	PYROCLASTIC	Gem Hill Formation—mainly tuff, tuff breccia, and tuffaceous sandstone (Miocene age uncertain) and Bobtail Quartz Latite Member—felsite breccia (Rosamond-Willow Springs area). Conejo Volcanics—agglomerate, pyroclastic and interbedded epiclastic rocks (Santa Monica Mts.). Rhyolite, andesite, and basalt breccia, agglomerate, scoriaceous flows, tuff, and tuff breccia (Santa Cruz Island).	
	MI	LOWER MIOCENE MARINE SEDIMENTARY ROCKS	"Tremble" Sandstone—arkosic sandstone, pebbly in part, calcareous in part (Santa Ynez River area); Rincon Shale—blue-gray, silty, micaceous shale (Santa Barbara-Ojai area); Vaqueros Formation—sandstone, siltstone, some conglomerate; in the Calliente Range divided into: Painted Rock Sandstone Member; Soda Lake Shale Member; Soda Lake Sandstone Member. Peto Formation—sandstone and interbedded clay shale (in part Oligocene). Unnamed sandstone and conglomerate on Santa Cruz Island, including glauconitic schist-bearing breccias and conglomerates "San Onofre Breccia" of Rand, 1931, ref. no. 80 (similar breccias occur on Anacapa Island where shown as M ^v). Unnamed brown to white sandstone, gray claystone and siltstone, in part middle Miocene (San Rafael Mts. area).	
	OC	OLIGOCENE NONMARINE SEDIMENTARY ROCKS	Sespe Formation—red to gray-green shale, sandstone, conglomerate, and breccias (probably ranges in age from late Eocene to early Miocene). Vasquez Formation—red to light-brown arkosic sandstone and conglomerate, some interbedded mafic siltstone and breccias (Oligocene or early Miocene age). Simmler Formation (Plush Ranch Formation of Carman)—red, green, and bluish-gray clay, arkosic sandstone, conglomerate, fanglomerate, and breccia (Oligocene or early Miocene age; Cuyama-Lockwood Valley areas). Tecuya Formation—red and green conglomerate sandstone and clay (early Miocene and Oligocene(?) age; south end San Joaquin Valley). Unnamed, greenish-gray to red conglomerate sandstone and mudstone of Miocene or Oligocene age (San Rafael Mountains).	
	O	OLIGOCENE MARINE SEDIMENTARY ROCKS	Caviota Formation—massive to thick-bedded, buff-weathering arkosic sandstone. "San Emigdio" Formation—alternate beds of yellowish and grayish sandstones and shales (San Emigdio Mts.).	
	Oligocene	OLIGOCENE VOLCANIC ROCKS:		
		OV	UNDIFFERENTIATED	Volcanic rocks of the Vasquez Formation, undifferentiated—black and reddish vesicular and amygdaloidal basalt and andesite, purple breccia, minor beds of tuff, associated with lacustrine strata in Tick Canyon containing borate beds. Volcanic rocks of the Vasquez Formation—olitic basalt, and andesite flows.
OV ^r		RHYOLITIC	Neenach Volcanic Formation—light-gray to pink rhyolitic felsite, Oligocene(?) and early or possibly middle Miocene.	
OV ^a		ANDESITIC	Volcanic rocks of the Vasquez Formation, andesitic—dark reddish-brown andesite flow breccias, including some greenish-white to buff tuff breccias. Neenach Volcanic Formation—dark-brown andesite, Oligocene(?) and early or possibly middle Miocene.	
OV ^b		BASALTIC	Volcanic rocks of the Vasquez Formation, basaltic—greenish-black basalts, commonly amygdaloidal. Volcanic rocks of the Plush Ranch Formation—black or red-brown hypersthene-augite-olivine basalt; some tuff and associated lacustrine strata that contain borate deposits. (Carman, ref. 12).	
E		Eocene MARINE SEDIMENTARY ROCKS	"Coldwater" Sandstone—arkosic sandstone, local oyster reefs, siltstone and shale; Sacate Formation—argillaceous to silty shale, some sandstone; Cozy Dell Shale—argillaceous to silty shale; Matijilla Sandstone—arkosic sandstone; Anita Shale—shale and some sandstone; Juncal Formation—shale and arkosic sandstone; Sierra Blanca Limestone—massive, hard, organic limestone (Santa Ynez, Topanga and San Rafael Mts.); Tejon Formation—sandstone, siltstone, shale, some conglomerate (San Emigdio-Tehachapi Mts.); Lajas Formation—conglomerate, sandstone, siltstone; Santa Susana Formation—concretionary shale, sandstone, conglomerate lenses (Paleocene in part). Unnamed rocks of middle or late Eocene age in Elmore Canyon. Unnamed Eocene siltstones, sandstones, shales and conglomerates in northern San Rafael Mountains, Santa Cruz Island, Piru Creek and Elizabeth Lake Canyon areas.	

STRATIGRAPHIC NOMENCLATURE—Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California.</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>Formally named formations grouped in sequence (separated by semicolons) are listed from youngest to oldest.</small>
PALEOCENE	Ep	PALEOCENE MARINE SEDIMENTARY ROCKS	Pattway Formation—siltstone, locally concretionary, sandstone, pebble-boulder conglomerate. "Martinez" Formation—sandstone, conglomerate, mudstone (includes some rocks of Eocene age); Las Virgenes Sandstone—gray sandstone (Simi Hills); Simi Conglomerate—well rounded pebbles and cobbles up to 1 foot diameter in matrix of coarse-grained arkosic sandstone; lenticular beds of sandstone. Unnamed sandstone interbedded with thin shale and conglomerate (Paleocene; Santa Cruz Island, Santa Monica Mts.). Unnamed conglomerates, sandstones, and siltstones (Paleocene and lower Eocene?; Castaic Creek-Elizabeth Canyon area).
	Tc	TERTIARY NONMARINE SEDIMENTARY ROCKS	Calliente Formation—calcareous, arkosic sandstone, red and gray mudstone, channel sand, and conglomerate; reddish-brown, thick-bedded, argillaceous, coarse-grained sandstone (now dated as early Miocene to middle Pliocene). Violin Breccia—of the San Gabriel fault scarp; grades into fine grained Ridge Route Formation to the east. (Miocene to Pliocene or Pleistocene age).
	Tm	TERTIARY MARINE SEDIMENTARY ROCKS	Unnamed shale and sandstone, some conglomerate, travertine, and tuff (except for travertine and arkose, has a marine aspect; age uncertain, may be Eocene, T. W. Dibble, Jr., written communication, 1968; located at intersection of Big Pine and San Andreas faults).
	Tv ^b	TERTIARY VOLCANIC ROCKS: BASALTIC	Basalt flows of the Calliente Formation—multiple olivine basalt flows, highly vesicular in part, includes breccias on some flow tops (Miocene age).
TERTIARY	TERTIARY INTRUSIVE (HYPABYSSAL) ROCKS:		
	Ti	UNDIFFERENTIATED	Intrusive rocks ranging from olivine basalt to dacite; middle Miocene(?) age (Santa Monica Mts.).
	Tif	RHYOLITIC	Bobtail Quartz Latite Member of Gem Hill Formation—intrusive facies including felsite and porphyritic facies, porphyry and perlite; Miocene(?) age; Rosamond-Willow Springs area. Rhyolite and obsidian plugs (Frazier Mountain area).
	Tia	ANDESITIC	Andesite porphyry of Miocene(?) or Oligocene(?) age (San Emigdio Mts.). Hornblende andesite sill (South Mountain).
	Tib	BASALTIC	Intrusive basalt and diabase of middle Miocene(?) age (Santa Monica Mts.).
	Ku	UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS	Ku: Jalama Formation—gray to black shale, hard, light-gray arkosic sandstone, and cobble conglomerate (Santa Ynez Mts.). "Chico" Formation—sandstone, shale and conglomerate. Unnamed light-gray sandstone dark-gray siltstone and shale, and minor cobble conglomerate.
	Kl	LOWER CRETACEOUS MARINE SEDIMENTARY ROCKS	Espada Formation—dark olive-gray shale, siltstone, and thin sandstone beds (San Rafael and Santa Ynez Mts.).
	Kjf	FRANCISCAN FORMATION	Franciscan Formation—sheared and severely deformed, dark greenish-gray gneiss, black claystone and chert (San Rafael and Santa Ynez Mts.).
	Kjfv	FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS	Greenstone of the Franciscan Formation and minor quantities of glaucophane schist (San Rafael and Santa Ynez Mts.).
	MESOZOIC	MESOZOIC GRANITIC ROCKS	
gr		UNDIFFERENTIATED	Granite and other quartz-bearing plutonic rocks (Soledad basin). Biotite granodiorite, quartz monzonite, and quartz diorite (Griffith Park). Areas of undivided, predominantly granitic rocks; locally may grade into rocks of banded "gneissic" structure (gneissic granite) or may be complexly associated with gneiss (Sawmill Mountain and Bouquet Reservoir areas). Mount Lowe "Granodiorite" and Parker "Quartz Diorite"—light-colored, foliated, quartz-poor, felspar-rich, porphyritic granitic rock of monzonitic to dioritic composition (Permian-Triassic; San Gabriel Mts.).
gr ^a		ADAMELLITE (QUARTZ MONZONITE), GRANITE, ALASKITE	Lebec Quartz Monzonite—fractured, gray hornblende-biotite quartz monzonite, with dark inclusions and pegmatites. Lebec Quartz Diorite—gray hornblende-biotite quartz monzonite. Mt. Pinos Granite—biotite granite and hornblende-biotite quartz diorite. Tejon Lookout Granite—biotite granite (Tehachapi Mts.). Unnamed granites and quartz monzonites (north of San Andreas fault).
gr ^b		GRANODIORITE	Unnamed granodiorite including some quartz diorite, quartz monzonite, and pink granite (San Gabriel Mts.).
gr ^c		TONALITE (QUARTZ DIORITE) AND DIORITE	Vermont Quartz Diorite—biotite quartz diorite (Griffith Park). Lar Quartz Diorite—biotite quartz diorite, commonly foliated (Griffith Park). Unnamed quartz diorites and diorite (Tehachapi Mts., San Gabriel-Verdugo Mts., Piru Creek area, San Emigdio Mts., Santa Cruz Island); includes migmatized metasedimentary rocks in Tehachapi Mts.
bi		MESOZOIC BASIC INTRUSIVE ROCKS	Hornblende diorite and gabbro (Tehachapi Mts., Verdugo Mts.).
ub		MESOZOIC ULTRABASIC INTRUSIVE ROCKS	Serpentine (San Rafael Mts.).
Ju		UPPER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Ju-Tr. Santa Monica Slate—highly foliated dark-gray to black slate, with minor amounts of mica schist, phyllite, and spotted cordierite slate (Late Jurassic pelecypods; formation probably Triassic in part, U.S. Geol. Survey Prof. Paper 420-A, p. 21-22).
PRE-CRETACEOUS METAMORPHIC ROCKS			
m		UNDIFFERENTIATED	Pelona Schist—highly foliated mica-chlorite-albite-quartz schists, some biotite schist, actinolite schist and quartzite, probably derived from sedimentary and pyroclastic rocks (age uncertain, generally considered to be Precambrian or Mesozoic). Diorite gneiss and some biotite schist (San Gabriel Mts.). Unnamed phyllite, chlorite schist and greenstone (Santa Cruz Island).
ls	ls = LIMESTONE AND/OR DOLOMITE	Thin strata of light blue-gray fine crystalline limestone in the Pelona Schist. Lenses of marble in granitic and metamorphic rocks (Tehachapi Mts. and area north of Mt. Pinos and Frazier Mtn.).	
ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS	Placerita Formation—graphite, sillimanite, and biotite schists, quartzite, and some limestone and dolomite (San Gabriel Mts.). Hornfels, quartzite, gneiss, marble, and schist (San Emigdio Mts.). Biotite gneiss, minor mica schist, quartzite, and hornfels (Tehachapi Mts.).	
mv	PRE-CRETACEOUS METAVOLCANIC ROCKS	Metamorphosed quartz latite (Soledad Mtn., Rosamond quadrangle).	
gr-m	PRE-CENOZOIC GRANITIC AND METAMORPHIC ROCKS	Diorite gneiss injected by granodiorite (migmatite) and in places includes the Placerita Formation (San Fernando quadrangle). Undifferentiated gneiss and granitic rocks (Mesozoic?) to Precambrian; Mint Canyon area.	
PALEOZOIC	PALEOZOIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS		
	IP	PALEOZOIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Bean Canyon Formation—dark-gray to black mica schist, phyllite, hornfels, and quartzite (age uncertain, may be Mesozoic; formation in part volcanic—see IP ^v ; Tehachapi Mts.).
	ls	ls = LIMESTONE AND/OR DOLOMITE	Limestone, dolomite and marble of the Bean Canyon Formation.
	IP ^v	PALEOZOIC METAVOLCANIC ROCKS	Metabasalt and metafelsite flows or sills of the Bean Canyon Formation (age uncertain, may be Mesozoic; Tehachapi Mts.).
	pCan	PRECAMBRIAN ANORTHOSITE	Anorthosite—Medium- to very coarse-grained; includes anorthosite intruded by Mesozoic(?) granite (San Gabriel Mts.).
	pCgr	UNDIVIDED PRECAMBRIAN GRANITIC ROCKS	Echo Granite—orange to pinkish-gray, locally foliated, quartz-rich granitic rocks (San Gabriel Mts.). Syenite phase of San Gabriel anorthosite complex (Mint Canyon area). Gabbro—"gabbroic and noritic rocks; altered rocks bordering anorthosite; ilmenite-magnetite gabbro and massive ilmenite-magnetite (San Gabriel Mts.).
PRECAMBRIAN	UNDIVIDED PRECAMBRIAN METAMORPHIC ROCKS:		
	pCg	GNEISS	Mendenhall Gneiss—"quartz-plagioclase gneiss with ferromagnesian minerals (San Gabriel Mts.). Gneiss, veined gneiss, banded gneiss and migmatites (Frazier Mountain and Mt. Pinos areas). Coarse and fine augen gneisses, layered quartzofeldspathic gneisses, minor amphibolites (Mint Canyon area). Banded amphibolite, hornfels, diorite and granite gneiss (Sawtooth Mts.). Foliated gneiss containing some recrystallized limestone and quartzite, local migmatite (Verdugo Mts.). For some of these rocks the attributed Precambrian age is uncertain. Complex of gneiss and lesser amounts of granitic rocks, locally includes migmatites and dark hornblende diorites (the granitic rocks are of magmatic origin or are recrystallized from gneiss and often grade into gneiss; age Mesozoic or older), Sawmill Mountain-Bouquet Reservoir area.
ls	ls = LIMESTONE AND/OR DOLOMITE	White coarsely crystalline marble lenses in gneiss, probably Precambrian, but possibly Paleozoic or Mesozoic (Sawmill Mountain area).	

NOTES

- Late Pliocene and early Pleistocene age. W. O. Addicott, 1965, U.S. Geol. Survey Prof. Paper 503-B, p. B-8. Radiometric dates (K-Ar) from ash layer(s) near base of Santa Barbara Formation range from 11.5 ± 2.4 m.y. to 7.8 ± 1.2 m.y., reported by Robert S. Yeats and W. A. McLaughlin, Geol. Soc. America Special Paper, in press, 1968.
- Probably Pleistocene, but may be late Pliocene in part.
- The Pico Formation includes rocks of early to late Pliocene age in the type locality in Ventura basin (Durham, D. L. and Yerkes, R. F., 1964, Geology and oil resources of the eastern Puente Hills area, southern California: U.S. Geol. Survey Prof. Paper 420-B, p. B 24-25); however, common usage in the U. S. A. basin restricts Pico to rocks of late Pliocene age. Modern U.S.G.S. usage assigns Pliocene marine rocks of the northeast L. A. basin to the upper member of the Fernando Formation.
- "Repetto" is defined and properly used only as a stage designation. The U.S. Geological Survey has abandoned the name "Repetto Formation" and assigns these rocks to the lower member of the Fernando Formation (Durham and Yerkes—see ref. footnote 4).
- Tequepis Sandstone and Sisquoc Formation are lateral equivalents; in part possibly of early Pliocene age in this map area.
- Also called Maricopa Shale in San Emigdio Mountains and Modelo Formation in Ventura basin. Ranges in age from middle Miocene (Relizian) to late Miocene (Mohnian). In Cuyama Valley area divided into Whiterock Bluff Shale Member, Santos Shale Member and Brant Canyon Formation.
- Oligocene or early Miocene age.
- Radiometric age date (Pb²⁰⁶/U²³⁸) of 245 ± 10 m.y., determined by L. T. Silver, 1968. Preliminary history for the crystalline complex of the central Transverse Ranges, Los Angeles County, California: Geol. Soc. Amer. Spec. Paper no. 101 (Abstracts), p. 20, 1967 and 1968.
- Isotopic age determinations (U-Th-Pb) on zircons from pegmatite and granophyre dikes in the western San Gabriel Mountains have yielded concordant ages of about 1200 million years. From structural and petrological relations, this is both a minimum and a probable age for the San Gabriel Mountains anorthosite complex. It is also a minimum age for the Mendenhall Gneiss . . . which has been intruded by the anorthosite complex. L. T. Silver, et al., 1963, Precambrian age determinations in the western San Gabriel Mountains, California: Jour. Geology, vol. 71, no. 2, p. 196-214.