DATE: 11 August 1984 SAMPLE:
CRUISE ID: SO 33 31DK box dredge
LOC: IV
AREA: C

	time (GMT)	LAT (N)	LON (W)	water depth (m)	wire
ON BOTTOM =	1256	6°16.8527	162°52.6657	13776	170
OFF BOTTOM =	1338	6°16.7332	162°52.6410	1378	144

time on bottom = 42 min

RECOVERY WEIGHT = total ~~300~~ kg crust <1 kg

CRUST THICKNESS = Veneer
1 generation

SUBSTRATE = bright red-brown glassy oxidized material. Unusual color
(Palagonite) (Blasenlava?)

COMMENTS:

* 1 microbio sample
* 1 bio sample

Sample # 1003: 3933 STDR I

Rock Type:	Basalt (AOB?) I	Basalt (AOB?) II
Color:	Medium brownish-gray (5YR 5/1)	Dark gray (N3)
Texture:	Subaphyric - phyric, hyalopilitic-intersertal	-do-
Phenocrysts:	Olivine <1 - ±5% (generally single euhedral phenocrysts)	-do-
Groundmass:	Olivine - Plagioclase - Pyroxene? - Glass - not determined	-do-
Vesicles:	Size: extremely variable (<<1mm - 8+mm)	<1 mm
	Shape: irregular	round
	%: extremely variable (<1% - 50±%)	<<1%
	Filling: empty - (usually) partial filling (locally) with calcite, zeolites, smectite?)	smectite
Alteration:	extensive (Oxidation [hematite?] smectitization, zeolitization, hydration) Olivine pxts completely altered to iddingsite	interstitial glass? looks fairly fresh -do-
Remarks:	Olivine basalt of diverse vesiculation and phenocryst density and distribution, but probably related to one flow unit, probably coupled subaerially and related to Basaltic Vent breccia II as the rocks bear many similarities, exposed surfaces bear thin (±1mm) Mn-oxide coatings, complete in protected areas, highly polished by water currents? on exposed surfaces.	Olivine basalt of consistently finer-grained texture than Basalt I and is much more dense, though appearance of groundmass and phenocrysts is much similar; probably a pahoehoe, less volatile-rich flow unit; exposed surfaces coated with thin (±1mm) sodose Mn-oxide crust.

3	4
Rock Type: Basalt (AOB?) III	Basalt (AOB?) IV
Color: Medium dark gray (N4)	Brownish-gray (5YR 3/1)
Texture: Sub-aphyric - intersertal	→ - do - perhaps hyalo-pilotaxitic
Phenocrysts: Olivine - <1%, ± 1 mm, single, euhedral	≠ ≠5%, ±1 mm dia, single euhedra
Groundmass: Plagioclase ? Olivine ? clinopyroxene ? not determined Glass ? Opaques ?	→ perhaps much more glass in matrix
Vesicles: %: ±5% Size: ± 1 mm Shape: irregular filling: calcite (50-75% amygdules)	— absent —
Alteration: moderate - extensive Olivine, both phenocrysts & groundmass are completely replaced; interstitial glass altered to smectite? much evidence for penetration by fluids along fractures (subsol. alteration)	extensive → - do - appears to be much more glass — totally altered to brown smectite - in the matrix
Remarks: Olivine basalt of similar aspect to Basalt II, but apparently more altered — may be the source of the unit exposed Surfaces coated [illegible] thin (<0.5 mm) zone of Mn-oxide coating which has been polished by bottom currents; apparently pebble-cobble-size components of a volcanic breccia, traces of which adhere to surfaces of the clasts	→ -do- even more altered probably the same unit as 31 DK 2

Rock Type: Olivine Basalt (AOB(?)) pillow b 5

Color: Medium brownish gray (5YR 5/1) to yellowish-brown (10YR 5/2)

Texture: Phyric hyalopilitic (?- intersertal

Phenocrysts: Olivine, ±10%, ±2 mm dia (single, euhedral)

Groundmass: Glass, Olivine, Plagioclase, Clinopyroxene?, Opaques? — not determined

Vesicles: absent

Alteration: Extensive (smectitization, oxidation, hydration, zeolitization(?)) Olivine phs & groundmass totally altered to pale green chlorophaeite; groundmass glass totally altered to smectite, etc.

Remarks: loose pebbles not related to other olivine basalt in the dredge or to the volcanogenic breccia; appear fragments of a pillow lava, some pieces bearing a ±5 mm-thick palagonite rind, appears to have undergone pre-eruptive, non-oxidizing hydration (olivine, glass) and post eruptive oxidative alteration (palagonitization, oxidation of groundmass glass)

AOB(?) Volcanogenic breccia 6

Yellowish-gray (5Y 7/2)

Clastic

(Matrix) predominant, AOB(?) rockfragments, ±45%.

(Cement) Calcite (calcareous ooze?) Smectite Zeolite? ±55%.

(Avge grain size) ±3 mm (fine gravel)

(Sorting) poor

(Grain shape) subangular - subrounded

(Porosity) ±25%.

(Permeability) ?20%

(Alteration) Smectitization zeolitization?

Remarks: rock made of lithoclasts derived from weathering and comminution of basaltic rocks and (perhaps) ash; looks like all rock fragments - perhaps shards for glass shards as well - not sufficient; grains range in size from silt size to pebble or cobble size and show evidence of considerable (subaerial) reworking before final deposition; thin coating (<1 of probable Mn-oxide on exposed surfaces

7

Rock Type: Basaltic Vent Breccia I

Color: Moderate reddish-brown (10R 4/6) to Grayish-Red (5R 4/2) to Dark grayish-red (5R 3/2)

Texture: Volcaniclastic, subaphyric, sub holohyaline, scoriaceous

Phenocrysts: olivine, <<1%, ±1mm

Groundmass: Glass ±99%

Vesicles: (extremely variable with distribution of flow lines) (also in size, shape) (<<1 mm — ±5mm dia)

Alteration: complete - nearly complete (Oxidation, sulfatization, hydration)

Remarks: Looks like a vent agglomerate/breccia-ish which easily hyaloclastic material piled up around a vent or vents, was incorporated in a later "Aa"-like flow of semi-liquid to semi-solid consistency. Perhaps the rusty brown color of the scoriaceous cementing glass can be explained by subsequent low-temperature hydrothermal alteration — especially since the rusty color is not completely homogeneous (in some areas it is light brown). much of the incorporated breccia pieces and blocks are also highly scoriaceous and are oxidized to grayish-red indicating oxidation at higher temperatures perhaps before being erupted - These blocks and pieces frequently bear large iddingsitized olivine phenocrysts; large clasts of the breccia display definite flow structure; the incorporated breccia pieces and blocks often display greenish nontronite & calcite deposits in vuggy areas.

exposed surfaces cooked with ±1 (± 1mm) layer of very shiny black crystalline MnO2 deposit

8

Basaltic Vent Breccia II

Color: Grayish-red (5R 4/2) to dark grayish-red (5R 3/2)

↔ more phyric (not much)

↔ - do -

↔ ±1%, ±2 mm dia

↔ - do -

↔ - do -

↔ - do -

↔ Similar to sample on left but lacking reddish-brown glass "cement"; in this case, far more clastic material is present and is more loosely bound together (more of an "Aa"-like flow perhaps). both these breccias were probably emplaced subaerially, possibly later altered by hydrothermal fluids subsequent to final deposition.

suberially emplaced probably