PROTOCOL DREDGE

Cruise No. SO 37	Stat. No. Equipm. 30 DKS	Location 3	Area I
Date 27.5.1985	Protocol written by : Herzig / Cremer		

Pinger on wire 3.00 m from dredge

	Time GMT	Latit. N	Longit. W	Water depth	Wire length	Tension (t)	Remarks
begin operation	0814	15°39.18	170°23.42	1273	—	.2	
begin lowering	0815	15°39.19	170°23.41	1273	—	.2	
+1. bottom contact	0930	15°39.59	170°23.62	1134	1520	2.8	
	0931	15°39.59	170°23.62	1138	1520	3.0	
	0934	15°39.57	170°23.61	1132	1530	2.9	
	0935	15°39.91	170°23.11	1130	1530	3.2	
	0944	15°39.87	170°23.05	1109	1540	4.3	
	0946	15°39.87	170°23.05	1111	1541	5.5	
	0950	15°39.86	170°23.04	1108	1570	3.3	
	0953	15°39.86	170°22.99	1104	1570	4.6	
	0956	15°39.86	170°22.94	1095	1571	4.0	
begin heaving	1003	15°39.84	170°22.86	1151	1515	3.8	
	1007	15°39.82	170°22.87	1153	1431	3.2	
	1008	15°39.81	170°22.87	1154	1403	4.0	
End Bottom Contact	1016	15°39.77	170°22.86	1224	1180	1.8	
Sampling	from ~1200		to	1080 m			
begin heaving	1000	15°39.85	170°22.86	1151	1562	2.6	
end heaving	1051	15°40.20	170°22.40	—	—	—	
end operation							

+Corrected data 15° 39.93' N 170° 23.15' W

ship course ...080°T... calcul. mean water depth.....m

total time of bottom contact.....min ship velocity..1...kts

bottom topography ..... length of dredge section .....m

Result

120.kg Total 90.kg Substrate

....kg Nodules ....kg Sediment

Pings 300 m from wire

	Time (GMT)	Lat N	Lon W	Water depth	Wire length	Tension
Begin Operation	08:14	15°39'19"	170°23'42"	1273	—	0.2
Begin Lowering	08:15	15°39'19"	170°23'41"	1273	—	0.2
+1 bottom contact	0930	15°30'59"	170°23'82"	1134	1520	2.8
End Bottom Contact	10:16	15°30'77"	170°22'86"	1224	1190	1.9

Sampling from 1200 – 1080 m

Begin Hauling	1006	15°39'85"	170°22'86"	1151	1562	2.6

Ship Course 080° T

Ship Velocity 1 Kt

total dredge Wt 120 kg 80 kg substrate

total Wt crust recovery: 30 kg

100% Crust coverage

100% talus

2 generations, older dense, laminar, younger dendritic; no obvious separation (i.e. phosphorite)

Surface topography: Subdued bottryoidal

max. thickness ~ 4 cm
min. thickness ~ 3 mm
} ~2 cm average

2 photos

Biology: red Corals

Gen. Substrates
chalk
phosphorite / volcanogenic breccia

Rock Type: Hornblende(?) Diorite(?) ≠ Andesite(?) ≠ Trachyte(?)

Color: Dark greenish-gray (5GY4/1) ≠ Medium Brownish-gray (5YR5/1) ≠ Brownish Orange pink 3YR 6/2

Texture: Intergranular (microgabbroic) ≠ Intergranular

Phenocrysts: Hornblende(?) ±10%, 2mm dia (combined) Augite(?) ±4% Plagioclase(?) ±3%, 1-2mm dia ≠ Olivine(?) ±5%

≠ Mafics (augite? hornblende?) ±10% Plagioclase ±10%

Groundmass: not differentiated — Mafics ±42% ≤0.05mm; Felsics ±45%) ≠ Not determined (grainsize too small to resolve w/ handlens) → do-

Vesicles: none (dense) ≠ ±3% irregular vesicles, ~1mm dia usually filled w/ calcite ≠ absent (dense)

Alteration: incipient (oxidation, smectitization weathering rind peripheral to sample clasts; smectitization of some groundmass material(?) & plagioclase(?) pxts ≠ extensive (oxidation of many groundmass mafics; smectitization, zeolitization(?); FeOH deposited along numerous fractures; Olivine(?) is altered universally to greenish-white smectite. ≠ superficial (oxidation of some groundmass components, smectitization of some feldspars

Xenoliths: large subangular cobble-size clasts, edges somewhat rounded; has basically the same mineral assemblage as andesites(?) from other areas on this site (i.e. 2005-1). Only the grainsize is coarser; has numerous blocky, small amoeboids of much coarser diorite(?) included. may be a fragment of a dike or trap-rock. occurs as talus with thin patina of manganese oxide encrustation — do- — do-

≠ Small cobble-size, rounded talus clast w/ patina of phosphorite then dendritic ferromanganese encrustation; origin of cobble unknown but appears to have been subjected to subaerial weathering prior to incorporation in slope debris volcanogenic breccia, dense to suggest that it may have been injected as trap.

Sample No: 30 ~~30054~~

Rock Type:	andesitic? Hyaloclastite Breccia	→ do → Volcanogenic Breccia	→ do
Color:	Greenish-gray (5GY 6/1) to moderate brown (5YR 3/4)	≠ Light olive gray (5Y 5/2) (mean color)	≠ Moderate yellowish-brown to very pale orange (10YR 5/4 to 10YR 8/2) (matrix)
Texture:	Pyroclastic	≠ clastic	→ do
Framework Grains:	Andesitic? hyaloclasts ±45%, ±3 mm dia Andesitic? clasts ±5%, ±5 mm dia Hornblende? crystals ±10%, ± 2 mm dia Augite? crystals open space ±20%, ±1 mm dia	→ do, ±18%, also, trachytic clasts, ±5%, phosphoritic clasts ±5%; → do, ±30%; → do, } ±5% → do, } → do, ±8%;	≠ andesitic clasts only, ± 30%, ± 2 mm dia
Cement:	Zeolites (phillipsite?) ±20%	≠ Phosphorite ±35%	→ do, (±70% less dense phos.
Average grain-size:	± 2 mm	≠ ±1 mm	→ ±1 cm phosphatized
(range:	≤1 mm–1 cm)	→ (range: ≤1 mm – do – 1 cm)	(range: ~5 mm → ±2 cm)
Average grain shape:	Subangular	≠ Subround	≠ angular
Sorting:	moderately poor	≠ poor	≠ extremely poor
Porosity:	Very high	→ Very high	≠ Moderately high
Permeability:	High	→ High	≠ Very low
Alteration:	extensive (Oxidation, Smectitization, Zeolitization)	→ do; also phosphorization of original chalk cement	→ do

Remarks: Not abundantly retrieved with dredge material, but found as clasts w/ 1 cm ± thick generations of dendritic internal growth, subdued botryoidal external growth on ferromanganese crust; probably out crop and talus as well. Resembles other such deposits from Black Friday Ridge.

Apparently extensive, though (±3.5 cm) pattern thin, irregular thickness bedded deposit related to 30 DS4 (phosphorite cement instead of zeolite, more closed framework, more varied component matrix) is both underlain and capped by other Volcanogenic breccia units whose cement is phosphorite and the volcanic components are less abundant and more reduced in size; the whole of these three units (represented in this sample) is capped by ± 0.5 cm-thick, 2 generation, (lower laminar, upper dendritic) largely and subtly botryoidal externally, w/ finely nodose supertopography; the lowermost unit (30 DS5) in this stratigraphic cross-section displays faint poorly-defined bedding; occurs as both outcrop & talus.

Flat slab of talus encrusted w/ ferromanganese deposit, a large cobble-size in dimension; contains slabby, prismatic clasts of andesite set in partially-phosphorized chalk. Some of the slabby clasts bear ±1 mm-thick crusts on their shaded sides. The chalk-phosphorite cement is penetrated by tabular dendrites; may represent a dislodged, disaggregated fragment of bedded unit.

ock Type:	Andesite? Volcanogenic Breccia	Andesite? Hyaloclastite Breccia	→ -do-
olor:	Olive gray (5Y 4/1) to Grayish-orange	Greenish-gray (5GY 6/1) to moderate brown (5YR 3/4)	→ Olive gray (5Y 4/1) to very pale yellowish-orange (10YR)
ze:	clastic →	pyroclastic	≠ clastic
ramework rains:	Andesite(?) clasts ± 10%, ± 1 cm dia Trachyte clasts ± 10%, ± 1 cm dia	→ -do-, but ± 50%, upto 3 cm dia. Andesitic hyaloclasts, ± 40%, upto 2 cm dia	→ -do-, but, ± 20%, ± 5 mm dia. avg; → -do-, but ± 25%, ± 5 mm dia avge & phosphorite clasts, ± 10%, ± 5 mm dia avge
ment:	phosphorized foraminiferal chalk, ± 80%	Zeolites ± 10% (Phillipsite?)	≠ phosphorite, ± 35%
erage grainsize:	± 1 cm (matrix)	± 1 cm	± 0.5 cm
nge:	± 2 mm – ± 1.5 cm; < 0.2 mm (cement)	(range: ≤ 0.5 mm – ± 3 cm)	(range: ± 1 mm – 2.5 cm)
erage grainshape:	subangular →	- do -	≠ angular, but a number are subrounded
ting:	extremely poor →	- do -	→ -do-
osity:	Very high	≠ High	≠ Moderately high
meability:	Very low	≠ High	≠ Very low
eration:	penultimate (Oxidation of some ferromagnesian matrix clasts, groundmass zeolitization, smectitization, phosphorization.	≠ variable – some pieces of andesitic material are only superficially smectitized, zeolitized; others are intensely so, and oxidized as well	≠ penultimate (Smectitization, zeolitization, oxidation of matrix components, phosphorization of cement)
marks:	See 30 DS 5	Probably a variant of 30 DS 4 but is composed of very large (pebble size) subangular-to-subrounded clasts included in andesitic hyaloclastite and forming rounded, flattened cobble-size, ferromanganese-encrusted (see 30 DS 4) talus clasts; (also available as outcrop); the clasts in this breccia, however are often heavy, scoriaceous andesitic material	Similar to other V.B.'s from this dredge but contains a wider range of volcanic/clastic components; framework components seem as a general sentiment to have been subjected to fartless subaerial water-working than many other of the V.B.'s from "Black Friday Ridge"; occurs as both outcrop and talus, and is coated with ferromanganese oxide deposit as in 30 DS 4.

mple No: S03730DS10

k Type: Andesite(?)/Phosphorite Volcanogenic Breccia.

or: Moderate yellowish-brown (10YR 5/4)

ture: clastic

mework rains: pebbles of other phosphorized V.B.'s ± 25%, ± 2 cm dia
phosphorite intraclasts ± 10%, ± 3 mm dia.
andesitic hyaloclasts ± 5%, ± 2 mm dia
" rock fragments ± 5%, ± 2 mm dia

ment: dense phosphorite ± 55%

in size: ± 2 mm dia.

ge grainshape: Subangular

ting: Poorly sorted

osity: Very low

eability: Extremely low

ration: Total (phosphatization of nearly everything; zeolitization?, Smectitization, Oxidation of matrix components)

Remarks: cobble-size talus clasts as pieces of outcrop seem; covered by thick ± generation laminar/dendritic ferromanganese oxide crust, as in (30 DS 4 Strange inclusion of pebble-size subrounded pieces composed of other, similar phosphorite breccia; main "cement" of this lithology displays a high degree of stratification, is very hard, heavy and dense though for all the advanced state of phosphorization, ± form "ghosts" and pellet?) "ghosts" can still be observed in abundance